

Monochromatic sumsets for colourings of \mathbb{R}

Dániel T. Soukup

<http://www.logic.univie.ac.at/~soukupd73/>



universität
wien

Supported in part by **FWF Grant I1921**.

Consistently, modulo some large cardinal,

if $f : \mathbb{R} \rightarrow r$ with $r \in \omega$ then there is an **infinite** $X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

$X + X = \{x + y : x, y \in X\}$ i.e. **repetitions are allowed.**

- How does this fit into the theory of partition relations?
- What goes into the proof of this result?
- Joint result with P. Komjáth, I. Leader, P. Russell, S. Shelah, D. T. Soukup, and Z. Vidnyánszky.

Consistently, modulo some large cardinal,

if $f : \mathbb{R} \rightarrow r$ with $r \in \omega$ then there is an **infinite** $X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

$X + X = \{x + y : x, y \in X\}$ i.e. **repetitions are allowed.**

- How does this fit into the theory of partition relations?
- What goes into the proof of this result?
- Joint result with P. Komjáth, I. Leader, P. Russell, S. Shelah, D. T. Soukup, and Z. Vidnyánszky.

Consistently, modulo some large cardinal,

if $f : \mathbb{R} \rightarrow r$ with $r \in \omega$ then there is an **infinite** $X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

$X + X = \{x + y : x, y \in X\}$ i.e. **repetitions are allowed.**

- How does this fit into the theory of partition relations?
- What goes into the proof of this result?
- Joint result with P. Komjáth, I. Leader, P. Russell, S. Shelah, D. T. Soukup, and Z. Vidnyánszky.

Consistently, modulo some large cardinal,

if $f : \mathbb{R} \rightarrow r$ with $r \in \omega$ then there is an **infinite** $X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

$X + X = \{x + y : x, y \in X\}$ i.e. **repetitions are allowed.**

- How does this fit into the theory of partition relations?
- What goes into the proof of this result?
- Joint result with P. Komjáth, I. Leader, P. Russell, S. Shelah, D. T. Soukup, and Z. Vidnyánszky.

Consistently, modulo some large cardinal,

if $f : \mathbb{R} \rightarrow r$ with $r \in \omega$ then there is an **infinite** $X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

$X + X = \{x + y : x, y \in X\}$ i.e. **repetitions are allowed.**

- How does this fit into the theory of partition relations?
- What goes into the proof of this result?
- Joint result with P. Komjáth, I. Leader, P. Russell, S. Shelah, D. T. Soukup, and Z. Vidnyánszky.

Evolving partition relations

... an incomplete overview ...

Evolving partition relations

If $f : \omega \rightarrow r$ then there is an infinite $X \subset \omega$ with $f \upharpoonright X$ constant.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

Evolving partition relations

If $f : [\omega]^k \rightarrow r$ then there is an infinite $X \subset \omega$ with $f \upharpoonright [X]^k$ constant.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

F. P. Ramsey, 1930

$$\omega \rightarrow (\omega)_r^k$$

Evolving partition relations

There is $f : [2^{\aleph_0}]^2 \rightarrow 2$ so that $f''[X]^2 = 2$ for any uncountable $X \subset 2^{\aleph_0}$.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

F. P. Ramsey, 1930

$$\omega \rightarrow (\omega)_r^k$$

W. Sierpinski, 1933

$$2^{\aleph_0} \not\rightarrow (\aleph_1)_2^2$$

Evolving partition relations

If $f : [\aleph_{k-1}^+]^k \rightarrow r$ then $f \upharpoonright [W]^k$ is constant for some uncountable $W \subseteq \aleph_{k-1}^+$.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

F. P. Ramsey, 1930

$$\omega \rightarrow (\omega)_r^k$$

W. Sierpinski, 1933

$$2^{\aleph_0} \not\rightarrow (\aleph_1)_2^2$$

Erdős, Rado 1956

$$\aleph_{k-1}^+ \rightarrow (\omega_1)_r^k \text{ for all } r < \omega.$$

Evolving partition relations

$FinSum(X) = \{x_0 + x_1 + \dots + x_\ell : x_0 < \dots < x_\ell \in X\}$ i.e. no repetitions.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

F. P. Ramsey, 1930

$$\omega \rightarrow (\omega)_r^k$$

W. Sierpinski, 1933

$$2^{\aleph_0} \not\rightarrow (\aleph_1)_2^2$$

Erdős, Rado 1956

$$\beth_{k-1}^+ \rightarrow (\omega_1)_r^k \text{ for all } r < \omega.$$

N. Hindman, 1974

if $f : \mathbb{N} \rightarrow r$ then there is some infinite $X \subseteq \mathbb{N}$
so that $f \upharpoonright FinSum(X)$ is constant.

Evolving partition relations

There is $f : [\aleph_1]^2 \rightarrow \aleph_1$ so that $f''[X]^2 = \aleph_1$ for any uncountable $X \subset \aleph_1$.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

F. P. Ramsey, 1930

$$\omega \rightarrow (\omega)_r^k$$

W. Sierpinski, 1933

$$2^{\aleph_0} \not\rightarrow (\aleph_1)_2^2$$

Erdős, Rado 1956

$$\beth_{k-1}^+ \rightarrow (\omega_1)_r^k \text{ for all } r < \omega.$$

N. Hindman, 1974

if $f : \mathbb{N} \rightarrow r$ then there is some infinite $X \subseteq \mathbb{N}$
so that $f \upharpoonright \text{FinSum}(X)$ is constant.

S. Todorcevic, 1987

$$\aleph_1 \not\rightarrow [\aleph_1]_{\aleph_1}^2$$

Evolving partition relations

If $f : [2^{\aleph_0}]^2 \rightarrow 3$ then there is an uncountable $X \subset 2^{\aleph_0}$ with $|f''[X]^2| \leq 2$.

P. H. Principle

$$\omega \rightarrow (\omega)_r^1$$

F. P. Ramsey, 1930

$$\omega \rightarrow (\omega)_r^k$$

W. Sierpinski, 1933

$$2^{\aleph_0} \not\rightarrow (\aleph_1)_2^2$$

Erdős, Rado 1956

$$\beth_{k-1}^+ \rightarrow (\omega_1)_r^k \text{ for all } r < \omega.$$

N. Hindman, 1974

if $f : \mathbb{N} \rightarrow r$ then there is some infinite $X \subseteq \mathbb{N}$
so that $f \upharpoonright \text{FinSum}(X)$ is constant.

S. Todorcevic, 1987

$$\aleph_1 \not\rightarrow [\aleph_1]_{\aleph_1}^2$$

S. Shelah, 1988

$$\text{Con}(2^{\aleph_0} \rightarrow [\aleph_1]_3^2)$$

Monochromatic sumsets in \mathbb{N}

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. **repetitions are not allowed**.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subseteq \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we **allow repetition**?

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. repetitions are not allowed.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subseteq \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we allow repetition?

Monochromatic sumsets in \mathbb{N}

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. **repetitions are not allowed**.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subseteq \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we **allow repetition**?

Monochromatic sumsets in \mathbb{N}

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. **repetitions are not allowed**.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subseteq \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we **allow repetition**?

Monochromatic sumsets in \mathbb{N}

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. **repetitions are not allowed**.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subseteq \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we **allow repetition**?

Monochromatic sumsets in \mathbb{N}

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. **repetitions are not allowed**.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subset \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we **allow repetition**?

Easy Ramsey consequence: if $f : \mathbb{N} \rightarrow r$ with $r \in \omega$ then there is an infinite $X \subseteq \mathbb{N}$ so that

$f \upharpoonright X \oplus X$ is constant.

Here $X \oplus X = \{x + y : x \neq y \in X\}$ i.e. **repetitions are not allowed**.

Proof:

- if $f : \mathbb{N} \rightarrow r$ then let $g : [\mathbb{N}]^2 \rightarrow r$ defined by $g(x, y) = f(x + y)$,
- if $X \subset \mathbb{N}$ and $g \upharpoonright [X]^2$ is constant then $f \upharpoonright X \oplus X$ is constant too.

[Owings, Hindman 1970s] What happens if we **allow repetition**?

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \bmod 4.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \bmod 4$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \bmod 4$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \bmod 4.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \bmod 4$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \bmod 4$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \pmod{4}.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \pmod{4}$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \pmod{4}$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \bmod 4.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \bmod 4$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \bmod 4$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \bmod 4.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \bmod 4$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \bmod 4$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \bmod 4.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \bmod 4$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \bmod 4$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \pmod{4}.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \pmod{4}$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \pmod{4}$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{N} - with repetitions?

$$X + X = X \oplus X \cup \{2x : x \in X\}.$$

There is $f : \mathbb{N} \rightarrow 4$ without infinite monochromatic sumsets:

$$f(x) = \lfloor \log_{\sqrt{2}}(x) \rfloor \bmod 4.$$

- Suppose that $X \subseteq \mathbb{N}$ is infinite and take $y \ll x \in X$.
- $|\log_{\sqrt{2}}(x) - \log_{\sqrt{2}}(x + y)| < 1$,
- $|f(x) - f(x + y)| \leq 1 \bmod 4$.
- $f(2x) = \lfloor \log_{\sqrt{2}}(x) + 2 \rfloor = f(x) + 2 \bmod 4$ so $f(2x) \neq f(x + y)$.

Can we do this with 2 colours???

Monochromatic sumsets in \mathbb{R}

Started in [Hindman, Leader, Strauss 2015]

If $f : \mathbb{R} \rightarrow r$ is Baire/Lebesgue measurable then there is a perfect $\emptyset \neq X \subseteq \mathbb{R}$ so that

$$f \upharpoonright X + X \text{ is constant.}$$

Without definability?

There is an $f : \mathbb{R} \rightarrow 2$ so that

$$f''X \oplus X = 2 \text{ for every uncountable } X \subset \mathbb{R}.$$

- [HLS] using CH, [Komjáth, DTS, Weiss] in ZFC, and consistently the number of colours is best possible.

Monochromatic sumsets in \mathbb{R}

Started in **[Hindman, Leader, Strauss 2015]**

If $f : \mathbb{R} \rightarrow r$ is **Baire/Lebesgue measurable** then there is a **perfect** $\emptyset \neq X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

Without definability?

There is an $f : \mathbb{R} \rightarrow 2$ so that

$f \upharpoonright X \oplus X = 2$ for every uncountable $X \subseteq \mathbb{R}$.

- **[HLS]** using CH, **[Komjáth, DTS, Weiss]** in ZFC, and consistently the number of colours is best possible.

Monochromatic sumsets in \mathbb{R}

Started in [Hindman, Leader, Strauss 2015]

If $f : \mathbb{R} \rightarrow r$ is **Baire/Lebesgue measurable** then there is a **perfect** $\emptyset \neq X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

Without definability?

There is an $f : \mathbb{R} \rightarrow 2$ so that

$f \upharpoonright X \oplus X = 2$ for every uncountable $X \subseteq \mathbb{R}$.

- [HLS] using CH, [Komjáth, DTS, Weiss] in ZFC, and consistently the number of colours is best possible.

Monochromatic sumsets in \mathbb{R}

Started in [Hindman, Leader, Strauss 2015]

If $f : \mathbb{R} \rightarrow r$ is **Baire/Lebesgue measurable** then there is a **perfect** $\emptyset \neq X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

Without definability?

There is an $f : \mathbb{R} \rightarrow 2$ so that

$f''X \oplus X = 2$ for every uncountable $X \subset \mathbb{R}$.

- [HLS] using CH, [Komjáth, DTS, Weiss] in ZFC, and consistently the number of colours is best possible.

Monochromatic sumsets in \mathbb{R}

Started in [Hindman, Leader, Strauss 2015]

If $f : \mathbb{R} \rightarrow r$ is **Baire/Lebesgue measurable** then there is a **perfect** $\emptyset \neq X \subseteq \mathbb{R}$ so that

$f \upharpoonright X + X$ is constant.

Without definability?

There is an $f : \mathbb{R} \rightarrow 2$ so that

$f''X \oplus X = 2$ for every uncountable $X \subset \mathbb{R}$.

- [HLS] using CH, [Komjáth, DTS, Weiss] in ZFC, and consistently the number of colours is best possible.

Continued by [Fernandez-Breton, Rinot 2016]:

- how to realize more colours on sets of the form $FinSum(X)$ (no repetitions),
- general theorems on **uncountable, commutative, cancellative semigroups G** .

Bottom line: without definability,

Infinite sumsets are best possible on \mathbb{R} with repetition allowed.

Continued by [Fernandez-Breton, Rinot 2016]:

- how to realize more colours on sets of the form $FinSum(X)$ (no repetitions),
- general theorems on **uncountable, commutative, cancellative semigroups** G .

Bottom line: without definability,

Infinite sumsets are best possible on \mathbb{R} with repetition allowed.

Continued by **[Fernandez-Breton, Rinot 2016]**:

- how to realize more colours on sets of the form $FinSum(X)$ (no repetitions),
- general theorems on **uncountable, commutative, cancellative semigroups G** .

Bottom line: without definability,

Infinite sumsets are best possible on \mathbb{R} with repetition allowed.

Continued by **[Fernandez-Breton, Rinot 2016]**:

- how to realize more colours on sets of the form *FinSum*(X) (no repetitions),
- general theorems on **uncountable, commutative, cancellative semigroups** G .

Bottom line: without definability,

Infinite sumsets are best possible on \mathbb{R} with repetition allowed.

Continued by **[Fernandez-Breton, Rinot 2016]**:

- how to realize more colours on sets of the form *FinSum*(X) (no repetitions),
- general theorems on **uncountable, commutative, cancellative semigroups** G .

Bottom line: without definability,

Infinite sumsets are best possible on \mathbb{R} with repetition allowed.

Continued by **[Fernandez-Breton, Rinot 2016]**:

- how to realize more colours on sets of the form *FinSum*(X) (no repetitions),
- general theorems on **uncountable, commutative, cancellative semigroups** G .

Bottom line: without definability,

Infinite sumsets are best possible on \mathbb{R} with repetition allowed.

Monochromatic sumsets - with repetitions

Recall: $\exists f : \mathbb{N} \rightarrow 4$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset \mathbb{N}$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

Monochromatic sumsets - with repetitions

Recall: $\exists f : \mathbb{N} \rightarrow 4$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset \mathbb{N}$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

Monochromatic sumsets - with repetitions

If $f : G(\kappa) \rightarrow r$ then $f \upharpoonright X + X$ is constant for some infinite $X \subset G(\kappa)$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.
Notation:

$$G(\kappa) \xrightarrow{+} (\aleph_0)_r$$

Monochromatic sumsets - with repetitions

$\exists f : G(\kappa) \rightarrow r$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset G(\kappa)$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

Notation:

$$G(\kappa) \overset{+}{\not\rightarrow} (\aleph_0)_r \text{ e.g. } \mathbb{N} \overset{+}{\not\rightarrow} (\aleph_0)_4$$

Monochromatic sumsets - with repetitions

$\exists f : G(\kappa) \rightarrow r$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset G(\kappa)$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

[Hindman, Leader, Strauss]

-
-
-
-

Monochromatic sumsets - with repetitions

$\exists f : \mathbb{Q} \rightarrow 72$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset \mathbb{Q}$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

[Hindman, Leader, Strauss]

- $\mathbb{Q} \not\overset{+}{\rightarrow} (\aleph_0)_{72}$.

-
-
-

Monochromatic sumsets - with repetitions

$\exists f : G(m) \rightarrow \mathbb{Z}_2$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset G(m)$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

[Hindman, Leader, Strauss]

- $\mathbb{Q} \not\rightarrow^+ (\aleph_0)_{\mathbb{Z}_2}$.
- $G(m) \not\rightarrow^+ (\aleph_0)_{\mathbb{Z}_2}$ for $m < \omega$.
-
-

Monochromatic sumsets - with repetitions

$\exists f : G(\aleph_0) \rightarrow 144$ so that $f \upharpoonright X + X$ is **not constant** for an infinite $X \subset G(\aleph_0)$.

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

[Hindman, Leader, Strauss]

- $\mathbb{Q} \not\rightarrow^+ (\aleph_0)_{72}$.
- $G(m) \not\rightarrow^+ (\aleph_0)_{72}$ for $m < \omega$.
- $G(\aleph_0) \not\rightarrow^+ (\aleph_0)_{144}$
-

Monochromatic sumsets - with repetitions

$\exists f : G(\aleph_m) \rightarrow 2^m \cdot 144$ so that $f \upharpoonright X + X$ is **not constant** for an infinite X .

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

[Hindman, Leader, Strauss]

- $\mathbb{Q} \overset{+}{\not\rightarrow} (\aleph_0)_{72}$.
- $G(m) \overset{+}{\not\rightarrow} (\aleph_0)_{72}$ for $m < \omega$.
- $G(\aleph_0) \overset{+}{\not\rightarrow} (\aleph_0)_{144}$
- $G(\aleph_m) \overset{+}{\not\rightarrow} (\aleph_0)_{2^m \cdot 144}$ for $m < \omega$.

Monochromatic sumsets - with repetitions

$\exists f : \mathbb{R} \rightarrow r$ so that $f \upharpoonright X + X$ is **not constant** for an infinite X .

Let $G(\kappa) = \bigoplus_{\kappa} \mathbb{Q}$ i.e. $x : \kappa \rightarrow \mathbb{Q}$ with $|\text{supp}(x)| < \omega$. E.g. $G(2^{\aleph_0}) \approx \mathbb{R}$.

[Hindman, Leader, Strauss]

- $\mathbb{Q} \not\overset{+}{\rightarrow} (\aleph_0)_{72}$.
- $G(m) \not\overset{+}{\rightarrow} (\aleph_0)_{72}$ for $m < \omega$.
- $G(\aleph_0) \not\overset{+}{\rightarrow} (\aleph_0)_{144}$
- $G(\aleph_m) \not\overset{+}{\rightarrow} (\aleph_0)_{2^m \cdot 144}$ for $m < \omega$.

Corollary

If $2^{\aleph_0} < \aleph_\omega$ then

$$\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$$

for some $r < \omega$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let

$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let $c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$

If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let

$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let $c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$

If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let

$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let $c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$

If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let

$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let $c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$

If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let

$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$

If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$



Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.

If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let


$c_s : [\kappa]^{|s|} \rightarrow 2$ by

$$c_s(a) = c(s * a).$$



Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2} s_1 * a_i$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2}s_1 * a_i$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2}s_1 * a_i$.

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.

Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2}s_1 * a_i$.

$$c(2x_i) = c_{s_1}(a_i) = 0 = c_{s_2}(a_i \cup a_j) = c(x_i + x_j).$$

Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.


Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2}s_1 * a_i$.

$$c(2x_i) = c_{s_1}(a_i) = 0 = c_{s_2}(a_i \cup a_j) = c(x_i + x_j).$$



Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.


Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2}s_1 * a_i$.

$$c(2x_i) = c_{s_1}(a_i) = 0 = c_{s_2}(a_i \cup a_j) = c(x_i + x_j).$$



Positive relations through 'position invariance'

Given $s \in \mathbb{Q}^{<\omega}$ and $a \in [\kappa]^{|s|}$, let


$$x = s * a \in \bigoplus_{\kappa} \mathbb{Q}$$

by $\text{supp}(x) = a$ and $x(a(i)) = s(i)$.

Suppose that $c : \bigoplus_{\kappa} \mathbb{Q} \rightarrow 2$, and let

$c_s : [\kappa]^{|s|} \rightarrow 2$ by


$$c_s(a) = c(s * a).$$



If κ is large enough then there is a large $W \subseteq \kappa$ so that c_{s_i} are constant on W for $i = 0, 1, 2$.


Assume that c_{s_1} and c_{s_2} are both constant 0.

Let $\alpha < \beta < \gamma_0 < \gamma_1 < \dots \in W$.



Let $a_i = \{\alpha, \beta, \gamma_i\}$ and $x_i = \frac{1}{2}s_1 * a_i$.

$$c(2x_i) = c_{s_1}(a_i) = 0 = c_{s_2}(a_i \cup a_j) = c(x_i + x_j).$$



If c_{s_0}, c_{s_2} have the same constant then we need $tp(W) = \omega + \omega$.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the Erdős-Rado theorem.

[DTS, Vidnyánszky]

$\Rightarrow G(c^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using polarized partition relations instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with CH and $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćević, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the Erdős-Rado theorem.

[DTS, Vidnyánszky]

$\Rightarrow G(c^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using polarized partition relations instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with CH and $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćević, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the Erdős-Rado theorem.

[DTS, Vidnyánszky]

$\Rightarrow G(c^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using polarized partition relations instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with CH and $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćevic, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the **Erdős-Rado theorem**.

[DTS, Vidnyánszky]

$\Rightarrow G(c^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using **polarized partition relations** instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with **CH** and $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćević, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the **Erdős-Rado theorem**.

[DTS, Vidnyánszky]

$\Rightarrow G(\mathfrak{c}^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using **polarized partition relations** instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with **CH** and $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćević, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the **Erdős-Rado theorem**.

[DTS, Vidnyánszky]

$\Rightarrow G(\mathfrak{c}^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using **polarized partition relations** instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with **CH** and $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćevic, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the **Erdős-Rado theorem**.

[DTS, Vidnyánszky]

$\Rightarrow G(\mathfrak{c}^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using **polarized partition relations** instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with **CH and** $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćević, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

[Komjáth] and [Leader, Russell] independently

$\Rightarrow G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ where $\kappa = \beth_{2r-1}(\aleph_0)$,

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ under GCH.

In ZFC maybe???

- using the **Erdős-Rado theorem**.

[DTS, Vidnyánszky]

$\Rightarrow G(c^+) \overset{+}{\rightarrow} (\aleph_0)_2$,

- using **polarized partition relations** instead.

$\Rightarrow G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ consistently from a measurable,
with **CH and** $2^{\aleph_1} = \aleph_{\omega+1}$,

- using a version of [Todorćevic, di Prisco] polarized relation for \aleph_ω with σ -closed forcing.

Positive relations on \mathbb{R} - the main result

Recall: if $2^{\aleph_0} < \aleph_\omega$ then $\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$ for some $r < \omega$.

Consistently, modulo an ω_1 -Erdős cardinal,

$$\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r \text{ for any } r < \omega.$$

The main ingredients are

- the **position invariance** from previous proofs, but
- **polarized relations** under $MA_{\aleph_1}(Knaster)$, and
- **[S. Shelah]** Consistently, modulo an ω_1 -Erdős cardinal, if $f : [2^{\aleph_0}]^{<\omega} \rightarrow r$ then there is an uncountable X and $F : X \leftrightarrow 2^\omega$ so that $f(\bar{x})$ **only depends on the finite tree $F[\bar{x}]$** for $\bar{x} \in [X]^{<\omega}$.

Positive relations on \mathbb{R} - the main result

Recall: if $2^{\aleph_0} < \aleph_\omega$ then $\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$ for some $r < \omega$.

Consistently, modulo an ω_1 -Erdős cardinal,

$$\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r \text{ for any } r < \omega.$$

The main ingredients are

- the **position invariance** from previous proofs, but
- **polarized relations** under $MA_{\aleph_1}(Knaster)$, and
- **[S. Shelah]** Consistently, modulo an ω_1 -Erdős cardinal, if $f : [2^{\aleph_0}]^{<\omega} \rightarrow r$ then there is an uncountable X and $F : X \leftrightarrow 2^\omega$ so that $f(\bar{x})$ **only depends on the finite tree $F[\bar{x}]$** for $\bar{x} \in [X]^{<\omega}$.

Positive relations on \mathbb{R} - the main result

Recall: if $2^{\aleph_0} < \aleph_\omega$ then $\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$ for some $r < \omega$.

Consistently, modulo an ω_1 -Erdős cardinal,

$$\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r \text{ for any } r < \omega.$$

The main ingredients are

- the **position invariance** from previous proofs, but
- **polarized relations** under $MA_{\aleph_1}(Knaster)$, and
- **[S. Shelah]** Consistently, modulo an ω_1 -Erdős cardinal, if $f : [2^{\aleph_0}]^{<\omega} \rightarrow r$ then there is an uncountable X and $F : X \leftrightarrow 2^\omega$ so that $f(\bar{x})$ **only depends on the finite tree $F[\bar{x}]$** for $\bar{x} \in [X]^{<\omega}$.

Positive relations on \mathbb{R} - the main result

Recall: if $2^{\aleph_0} < \aleph_\omega$ then $\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$ for some $r < \omega$.

Consistently, modulo an ω_1 -Erdős cardinal,

$$\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r \text{ for any } r < \omega.$$

The main ingredients are

- the **position invariance** from previous proofs, but
- **polarized relations** under $MA_{\aleph_1}(Knaster)$, and
- **[S. Shelah]** Consistently, modulo an ω_1 -Erdős cardinal, if $f : [2^{\aleph_0}]^{<\omega} \rightarrow r$ then there is an uncountable X and $F : X \leftrightarrow 2^\omega$ so that $f(\bar{x})$ **only depends on the finite tree $F[\bar{x}]$** for $\bar{x} \in [X]^{<\omega}$.

Positive relations on \mathbb{R} - the main result

Recall: if $2^{\aleph_0} < \aleph_\omega$ then $\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$ for some $r < \omega$.

Consistently, modulo an ω_1 -Erdős cardinal,

$$\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r \text{ for any } r < \omega.$$

The main ingredients are

- the **position invariance** from previous proofs, but
- **polarized relations** under $MA_{\aleph_1}(\text{Knaster})$, and
- [S. Shelah] Consistently, modulo an ω_1 -Erdős cardinal, if $f : [2^{\aleph_0}]^{<\omega} \rightarrow r$ then there is an uncountable X and $F : X \leftrightarrow 2^\omega$ so that $f(\bar{x})$ **only depends on the finite tree $F[\bar{x}]$** for $\bar{x} \in [X]^{<\omega}$.

Positive relations on \mathbb{R} - the main result

Recall: if $2^{\aleph_0} < \aleph_\omega$ then $\mathbb{R} \not\overset{+}{\rightarrow} (\aleph_0)_r$ for some $r < \omega$.

Consistently, modulo an ω_1 -Erdős cardinal,

$$\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r \text{ for any } r < \omega.$$

The main ingredients are

- the **position invariance** from previous proofs, but
- **polarized relations** under $MA_{\aleph_1}(Knaster)$, and
- **[S. Shelah]** Consistently, modulo an ω_1 -Erdős cardinal, if $f : [2^{\aleph_0}]^{<\omega} \rightarrow r$ then there is an uncountable X and $F : X \leftrightarrow 2^\omega$ so that **$f(\bar{x})$ only depends on the finite tree $F[\bar{x}]$** for $\bar{x} \in [X]^{<\omega}$.

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_r) \not\rightarrow^+ (\aleph_0)_r$, for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$, if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$, without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$
for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_r) \not\rightarrow^+ (\aleph_0)_r$, for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$, if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$, without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \not\rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \not\rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$
for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \not\rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \dots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \not\rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \dots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \not\rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$
for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \dots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \dots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \dots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\rightarrow^+ (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \rightarrow^+ (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \rightarrow^+ (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\rightarrow^+ (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\overset{+}{\rightarrow} (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \overset{+}{\rightarrow} (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \dots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems

[Owings, 1974]

- $\mathbb{N} \not\overset{+}{\rightarrow} (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\overset{+}{\rightarrow} (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???

Various open problems – Thank you for your attention!

[Owings, 1974]

- $\mathbb{N} \not\overset{+}{\rightarrow} (\aleph_0)_2$???

Connected to our results:

- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_2$ in ZFC??
- $G(\aleph_\omega) \overset{+}{\rightarrow} (\aleph_0)_r$ for $r < \omega$ in ZFC???
- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r$ if 2^{\aleph_0} is real-valued measurable?
- $\mathbb{R} \overset{+}{\rightarrow} (\aleph_0)_r$ without large cardinals?
- unbalanced sumsets $X + Y$ (or $X \oplus Y$)?

What is the smallest r so that $G(\kappa) \not\overset{+}{\rightarrow} (\aleph_0)_r$ for a particular κ (finite, or \aleph_m)??

Monochromatic k -sumsets: $X + X + \cdots + X$?

[HLS] There is a finite colouring of $G(\aleph_n)$ with no infinite monochromatic k -sumsets ($n < \omega$),

[DS, Vidnyánszky] There is a finite coloring of \mathbb{R} with no infinite monochromatic k -sumsets for $k \geq 3$.

We are far from a complete picture.

[Shelah, 1988]

- Is $2^{\aleph_0} = \aleph_m \rightarrow [\aleph_1]_3^2$ consistent for some $m < \omega$???
- Is $2^{\aleph_0} > \lambda \rightarrow [\aleph_1]_3^2$ consistent???